

Sweden it is replaced by a related species, *Stigmella benanderella* (Wolff, 1955).

Acknowledgments

I wish to extend thanks to Dr. W. A. Foster for permission to inspect the collections at Cambridge University and to borrow the original specimen of *S. repentiella* for study; to Dr. J. D. Bradley who confirmed the determination by dissection; to the officers of the Kent Trust for Nature Conservation for information about the foodplant in Kent; to Dr. F. H. Perring for a map showing its distribution in the British Isles; to Dr. Neils L. Wolff for information about the foodplant and status of the insect in Denmark and elsewhere on the Continent; to Mr. S. N. A. Jacobs for excellent drawings of the imago and larval mine; and particularly to the Sandwich Bay bird-watchers who not only uncomplainingly tolerated entomologists who intruded on their preserves and impeded their activities, but have also cleared away some of the intrusive sea buckthorn (*Hippophaë rhamnoides*) to give the creeping willow a better chance to flourish.

References

- Clapham, A. R., Tutin, T. G. and Warburg, E. F., 1962. *Flora of the British Isles* (2nd Ed.). Cambridge.
 Hering, E. M., 1951. *Biology of the leaf miners*. 'S-Gravenhage.
 Tutin, E. G. et al., 1972-76. *Flora Europaea*. Cambridge.
 Wolff, N. L., 1955. *Nepticula repentiella* sp.n. (Lepidoptera Nepticulidae). *Ent. Meddr.*, 27: 82.
 Wolff, N. L., 1955. *Nepticula benanderella* n.sp. (Lep. Nepticulidae). *Opusc. Ent.*, 20: 49.

POLYDRUSUS SERICEUS SCHALL. (COL.: CURCULIONIDAE) IN SUSSEX. — Whilst working for the new British weevil *Magdalis memnonia* Gyll. in company with its discoverer, Mr. P. J. Hodge, in Friston Forest, East Sussex, on 15th June, 1974 (cf. Allen, 1975, *Ent. mon. Mag.*, 111: 96), I took a single example of this middle-sized green *Polydrusus* by sweeping in a ride. Mr. Hodge had never before seen the insect, but the following year found it in profusion at the same place on young oaks, etc.; and likewise in 1976 but less abundantly. *P. sericeus* is a particularly local weevil in this country, with its headquarters in North Hants.: Kimpton, Thruxton, Andover, Harewood Forest — all four places being in the same area. At the first two it was taken last century; at the third, by the late Philip Harwood, and one by myself in 1933 in a wood on the downs; at the fourth, by Joy and Tomlin in numbers and as lately as 1974, still in plenty, by Mr. D. Appleton. There are, besides, a few other scattered southern records. In Kent it must be excessively local, its sole station known to me being a cherry tree in the late Dr. Masee's garden at East Malling; possibly he may have met with it elsewhere in the district, but I never heard that he did. — A. A. ALLEN.